

Post Chemotherapy ≤ 9 years of age	Post Chemotherapy ≥ 10 years of age	Post HSCT ≤ 9 years of age	Post HSCT ≥ 10 years of age	Glossary
<p>Booster doses for children who have received their primary course of vaccination before diagnosis.</p> <p>Please Note: These guidelines apply to children who are < 2 yrs post treatment</p> <p>Booster dose/s commence 6 months after chemotherapy and if underlying illness is in remission.</p> <ul style="list-style-type: none"> • Infanrix Hexa • Prevenar 13 • Meningococcal ACWY • Pneumovax 23 <ul style="list-style-type: none"> ○ minimum 8 weeks post Prevenar 13 ○ consider if ≥ 2 years of age 	<p>Booster doses for children who have received their primary course of vaccination before diagnosis.</p> <p>Please Note: These guidelines apply to children who are < 2 yrs post treatment</p> <p>Booster dose/s commence 6 months after chemotherapy and if underlying illness is in remission.</p> <ul style="list-style-type: none"> • Boostrix-IPV • Prevenar 13 • Meningococcal ACWY • Hepatitis B <ul style="list-style-type: none"> ○ Paediatric dose ≤ 10 years ○ Adult dose ≥ 11 years • Gardasil (Children ≥ 12 yrs. of age) <ul style="list-style-type: none"> ○ Single dose if completed primary course ○ Recommence schedule if partially completed prior to treatment ○ If no previous 4vHPV vaccine require 3-doses: 0, 2 and 6 months • Pneumovax 23 <ul style="list-style-type: none"> ○ minimum 8 weeks post Prevenar 13 	<p>Please Note: These guidelines apply to children who are < 2 yrs post treatment</p> <p>6 Months post HSCT (Visit 1)</p> <ul style="list-style-type: none"> • Prevenar 13 – dose 1 • Infanrix Hexa – dose 1 • Meningococcal ACWY – dose 1 • Bexsero - dose 1 <p>Visit 2 (8 weeks post visit 1)</p> <ul style="list-style-type: none"> • Prevenar 13 – dose 2 • Infanrix Hexa – dose 2 • Meningococcal ACWY – dose 2 • Bexsero – dose 2 <p>Visit 3 (8 weeks post visit 2)</p> <ul style="list-style-type: none"> • Prevenar 13 – dose 3 • Infanrix Hexa – dose 3 <p>Visit 4 (8 weeks post visit 3)</p> <ul style="list-style-type: none"> • Pneumovax 23 	<p>Please Note: These guidelines apply to children who are < 2 yrs post treatment</p> <p>6 Months post HSCT (Visit 1)</p> <ul style="list-style-type: none"> • Prevenar 13 – dose 1 • Boostrix-IPV – dose 1 • Menitorix – dose 1 • Hepatitis B (Adult) – dose 1 • Meningococcal ACWY – dose 1 • Bexsero – dose 1 • Gardasil - dose 1 (≥ 12 yrs of age) <p>Visit 2 (8 weeks post visit 1)</p> <ul style="list-style-type: none"> • Prevenar 13 – dose 2 • Boostrix-IPV – dose 2 • Menitorix – dose 2 • Hepatitis B (Adult) – dose 2 • Meningococcal ACWY – dose 2 • Bexsero – dose 2 • Gardasil - dose 2 (≥ 12 yrs of age) <p>Visit 3 (8 weeks post visit 2)</p> <ul style="list-style-type: none"> • Prevenar 13 – dose 3 • Boostrix-IPV – dose 3 • Menitorix – dose 3 • Hepatitis B (Adult) – Dose 3 <p>Visit 4 (8 weeks post visit 3)</p> <ul style="list-style-type: none"> • Pneumovax 23 <p>Gardasil – dose 3 (≥ 12 yrs of age)</p>	<p>Boostrix-IPV</p> <ul style="list-style-type: none"> * Diphtheria * Tetanus * Pertussis * Inactivated Polio <p>Gardasil</p> <ul style="list-style-type: none"> * Human Papillomavirus <p>Hepatitis B</p> <ul style="list-style-type: none"> * adult dose or paediatric dose <p>Menitorix</p> <ul style="list-style-type: none"> * Haemophilus influenza type b * Meningococcal C <p>Infanrix Hexa</p> <ul style="list-style-type: none"> * Diphtheria * Tetanus * Pertussis * Polio * Hib * Hepatitis B (paed) <p>Meningococcal ACWY</p> <ul style="list-style-type: none"> * Menveo * Menactra * Nimenrix <p>Pneumovax 23</p> <ul style="list-style-type: none"> * Pneumococcal Polysaccharide <p>Prevenar 13</p> <ul style="list-style-type: none"> * Pneumococcal Conjugate
<p>Post Chemotherapy Live vaccines</p>	<p>Post Chemotherapy Live vaccines</p>	<p>Post HSCT Live vaccines</p>	<p>Post HSCT Live vaccines</p>	
<p>Only give on consultation with oncologist</p> <ul style="list-style-type: none"> • Measles, Mumps and Rubella 2 Doses • Varicella 2 Doses <p>Please note</p> <ul style="list-style-type: none"> • Live vaccines must be given on the same day or 4 weeks apart • Do not use combination Measles, Mumps and Rubella (MMRV) vaccine in children < 4 yrs of age for the 1st measles vaccine • If child has previously received MMR Pre Chemo. MMRV <u>can</u> be used as dose 1 for Post Chemo • Combination MMRV may be used as the 2nd dose for all 	<p>Only give on consultation with oncologist</p> <ul style="list-style-type: none"> • Measles, Mumps and Rubella 2 doses • Varicella 2 doses <p>Please note</p> <ul style="list-style-type: none"> • <i>Combination MMRV vaccine may be used in this group for both doses</i> 	<p>Only give on consultation with oncologist</p> <ul style="list-style-type: none"> • Measles, Mumps and Rubella 2 doses • Varicella 2 doses <p>Please note</p> <ul style="list-style-type: none"> • Live vaccines must be given on the same day or 4 weeks apart • Do not use combination MMRV vaccine in children < 4 yrs of age for the 1st measles vaccine unless they have previously received an MMR prior to HSCT • Combination MMRV may be used as the 2nd dose for all 	<p>Only give on consultation with oncologist</p> <ul style="list-style-type: none"> • Measles, Mumps and Rubella 2 doses • Varicella 2 doses <p>Please note</p> <ul style="list-style-type: none"> • <i>Combination MMRV vaccine may be used in this group for both doses</i> 	<p>Measles, Mumps and Rubella</p> <ul style="list-style-type: none"> * Priorix * MMRII <p>MMRV vaccine</p> <ul style="list-style-type: none"> * Priorix -tetra * ProQuad <p>Varicella</p> <ul style="list-style-type: none"> * Varivax * Varilrix <p>Bexsero</p> <ul style="list-style-type: none"> * Meningococcal B
<p>Influenza vaccine</p> <p>Recommended annually if ≥ 6 months. Require 2 doses in the first year of receiving the vaccine regardless of age (4 weeks apart) http://www.mvec.vic.edu.au/immunisation-references/influenza-vaccine-recommendations/</p> <p>If splenectomy/hyposplenism refer to MVEC for asplenia guideline www.mvec.vic.edu.au/immunisation-references/aspleniahyposplenial/</p>				<p>Note: Menitorix and Menveo can be given on the same day or 4 weeks apart</p>